


5 kg



5 kg

5 kg

Intubation – prepare ONE size tube above and below recommended size			
ETT size – mm – CUFFED	3 mm	NG tube	6 Fr
ETT size – mm – UNCUFFED	3.5 mm	ICC tube	8 - 12 Fr
ETT at lips – cm	10 cm	LMA	1
ETT at nose – cm	12 cm	IDC	6 Fr

ANAPHYLAXIS		
IM Adrenaline (Epinephrine) 1:1000 (1 mg/mL)		
Dose	Volume	Autoinjector
50 microg	0.05 mL	150 microg

*Use autoinjector only if adrenaline 1:1000 not available

Resuscitation	Vial concentration	Recommended dose/kg	Preparation		Dose	Final volume to administer	Administration	
			Dilution – Sodium Chloride 0.9%	Final concentration				
Adrenaline (Epinephrine) 1:10 000 (1 mg/10 mL)	100 microg/mL	10 microg/kg	Undiluted		100 microg/mL	50 microg	0.5 mL	Push
DC shock – VF/ pulseless VT		4 Joule/kg	Round up energy level to next highest setting on defibrillator			20 Joule		Use infant or paediatric pads
AmiODAROne (150 mg/3 mL)	50 mg/mL	5 mg/kg	<i>Dilute 3 mL (150 mg) to 15 mL in glucose 5%</i>		10 mg/mL	25 mg	2.5 mL	Push over 5 mins
Fluid Bolus		10 mL/kg	Sodium Chloride 0.9%				50 mL	Push
Fluid Bolus		20 mL/kg	Sodium Chloride 0.9%				100 mL	Push
Glucose 10%	100 mg/mL	2 mL/kg	Glucose 10%		100 mg/mL		10 mL	Push
Adenosine (6 mg/2 mL) – 1st dose	3 mg/mL	0.1 mg/kg	Undiluted		3 mg/mL	0.5 mg	0.17 mL	Push via proximal vein or CVL – Follow immediately by a 10 - 20 mL fast flush
Adenosine (6 mg/2 mL) – 2nd dose	3 mg/mL	0.2 mg/kg				1 mg	0.33 mL	
Adenosine (6 mg/2 mL) – 3rd dose	3 mg/mL	0.3 mg/kg				1.5 mg	0.5 mL	
Synchronised Cardioversion		1 Joule/kg	Round up energy level to next highest setting on defibrillator			5 Joule		Use infant or paediatric pads
		2 Joule/kg				10 Joule		
Atropine (600 microg/mL)	600 microg/mL	20 microg/kg	Dilute 1 mL (600 microg) to 6 mL		100 microg/mL	100 microg	1 mL	Push
Push dose pressors – Doses may be repeated if required								
Adrenaline (Epinephrine) 1:10 000 (1 mg/10 mL)	100 microg/mL	1 microg/kg	Dilute 1 mL (100 microg) to 10 mL		10 microg/mL	5 microg	0.5 mL	Push
Metaraminol (Syringe 5 mg/10 mL)	500 microg/mL	10 microg/kg	Consider Adrenaline (Epinephrine) Push Dose Pressor		Consult	Consult	Consult	Push

Induction agents	Vial concentration	Recommended dose/kg	Dilution – Sodium Chloride 0.9%	Final concentration	Dose	Final volume	Administration
Fentanyl (100 microg/2 mL)	50 microg/mL	2-5 microg/kg	Dilute 2 mL (100 microg) to 10 mL	10 microg/mL	10 microg	1 mL	Push over 1 - 3 mins
Ketamine (200 mg/2 mL)	100 mg/mL	1 - 2 mg/kg	Dilute 2 mL (200 mg) to 20 mL	10 mg/mL	5 mg	0.5 mL	Push over 60 secs
PropOFol (200 mg/20 mL)	10 mg/mL	2 - 3 mg/kg	Undiluted	10 mg/mL	10 mg	1 mL	Push over 30 secs
Midazolam	Various strengths	0.1 - 0.2 mg/kg	Dilute to 1 mg/mL regardless of ampoule strength	1 mg/mL	0.5 mg	0.5 mL	Push over 2 - 3 mins

Paralytic agents	Vial concentration	Recommended dose/kg	Dilution – Sodium Chloride 0.9%	Final concentration	Dose	Final volume	Administration
Rocuronium (50 mg/5 mL)	10 mg/mL	1.2 mg/kg	Undiluted	10 mg/mL	6 mg	0.6 mL	Push
Suxamethonium (100 mg/2 mL)	50 mg/mL	2 mg/kg	Dilute 2 mL (100 mg) to 10 mL	10 mg/mL	10 mg	1 mL	Push
Vecuronium (10 mg)	10 mg	0.1 mg/kg	<i>Reconstitute vial with 10 mL WFI</i>	1 mg/mL	0.5 mg	0.5 mL	Push

5kg

Reversal agents	Vial concentration	Recommended dose/kg	Preparation		Dose	Final volume to administer	Administration
			Dilution – Sodium Chloride 0.9%	Final concentration			
Sugammadex (200 mg/2 mL) Rocuronium reversal	100 mg/mL	16 mg/kg	Undiluted	100 mg/mL	80 mg	0.8 mL	Push
Flumazenil (500 microg/5 mL) Benzodiazepine reversal	100 microg/mL	5 microg/kg	Undiluted	100 microg/mL	25 microg	0.25 mL	Push – Every 60 secs Max single dose 200 microg Max total dose 2000 microg
Naloxone (400 microg/mL) Opioid reversal	400 microg/mL	10 microg/kg	Undiluted	400 microg/mL	50 microg	0.13 mL	Push – Every 2 - 3 mins May be given IM

Respiratory	Vial concentration	Recommended dose/kg	Dilution – Sodium Chloride 0.9%	Final concentration	Dose	Final volume	Administration
Nebulised Adrenaline (Epinephrine) 1:1000	1 mg/mL		Undiluted	1 mg/mL	5 mg	5 mL	Via nebuliser
Dexamethasone (4 mg/mL)	4 mg/mL	0.3 mg/kg	Undiluted	4 mg/mL	1.5 mg	0.38 mL	IV or IM
Magnesium Sulfate (10 mmol/5 mL)	2 mmol/mL	0.2 mmol/kg	Dilute 5 mL (10 mmol) to 50 mL	0.2 mmol/mL	Consult	Consult	Infuse over 20 mins
Hydrocortisone (100 mg + 2 mL diluent)	50 mg/ mL	4 mg/kg	<i>Reconstitute vial with 2 mL WFI</i>	50 mg/mL	20 mg	0.4 mL	Push over 30 secs or IM
Methylprednisolone (40 mg/mL) sodium succinate	40 mg/mL	1 mg/kg	Dilute 1 mL (40 mg) to 4 mL	10 mg/mL	Consult	Consult	Push over 5 mins Sodium succinate ONLY
Salbutamol (5 mg/5 mL)	1 mg/mL	0.1 mg/kg	Dilute 5 mL (5 mg) to 50 mL	0.1 mg/mL	Consult	Consult	Load – Infuse over 20 mins
AminOPHYLLine (250 mg/10 mL)	25 mg/mL	5 mg/kg	Dilute 10 mL (250 mg) to 50 mL	5 mg/mL	Consult	Consult	Load – Infuse over 30 mins

Neurology/seizures	Vial concentration	Recommended dose/kg	Dilution – Sodium Chloride 0.9%	Final concentration	Dose	Final volume	Administration
Midazolam – IV	Various strengths	0.15 mg/kg	Dilute to 1 mg/mL regardless of ampoule strength	1 mg/mL	0.75 mg	0.75 mL	Push
Midazolam – IM	5 mg/mL	0.2 mg/kg	Undiluted	5 mg/mL	1 mg	0.2 mL	IM
Midazolam – Buccal/Nasal	5 mg/mL	0.3 mg/kg	Undiluted	5 mg/mL	1.5 mg	0.3 mL	Drip dose into alternate nostrils or inside cheek
Phenytoin (100 mg/2 mL) (250 mg/5 mL)	50 mg/mL	20 mg/kg	Dilute 2 mL (100 mg) to 10 mL	10 mg/mL	100 mg	10 mL	Infuse over 20 mins *use 0.22 micron filter*
Phenobarbitone (Phenobarbital) (200 mg/mL)	200 mg/mL	20 mg/kg	Dilute 1 mL (200 mg) to 10 mL	20 mg/mL	100 mg	5 mL	Infuse over 20 mins
Levetiracetam (500 mg/5 mL)	100 mg/mL	60 mg/kg	Dilute 5 mL (500 mg) to 10 mL	50 mg/mL	300 mg	6 mL	Push over 5 mins
Mannitol 20%	0.2 g/mL	0.5 g (2.5 mL)/kg	Pre-mixed bag	0.2 g/mL	2.5 g	12.5 mL	Infuse over 10 mins *use 5 micron filter*
Sodium Chloride 3% – Hypertonic *For raised ICP or hyponatremic seizures*	0.5 mmol/mL	3 mL/kg	Pre-mixed bag	0.5 mmol/mL	15 mL	15 mL	Infuse over 10 mins via central/large vein

5kg

Electrolytes	Vial concentration	Recommended dose/kg	Preparation		Dose	Final volume to administer	Administration
			Dilution – Sodium Chloride 0.9%	Final concentration			
Hypokalaemia (↓ Potassium) Potassium Chloride 10 mmol in 0.29% sodium chloride (100 mL)	0.1 mmol/mL	0.3 mmol/kg	Pre-mixed bag	0.1 mmol/mL	1.5 mmol	15 mL	Infuse over 1 hour
Hyperkalaemia (↑ Potassium) Calcium gluconate (2.2 mmol/10 mL)	0.22 mmol/mL	0.11 mmol/kg	Undiluted	0.22 mmol/mL	0.55 mmol	2.5 mL	Large vein push over 3 - 5 mins DO NOT give with sodium bicarbonate
Salbutamol Nebules	2.5 mg/2.5 mL	Age based	Dilute to 4 mL	–	2.5 mg	–	Inhale via nebuliser
Frusemide (Furosemide) (20 mg/2 mL)	10 mg/mL	1 mg/kg	Dilute 2 mL (20 mg) to 20 mL	1 mg/mL	5 mg	5 mL	Push over 5 mins
Glucose 10% (with insulin below)	See Infusion guide for doses and administration directions						
Insulin – Actrapid (300 units/3 mL)							
Sodium Bicarbonate 8.4%	1 mmol/mL	1 mmol/kg	Dilute 10 mL (10 mmol) to 20 mL	0.5 mmol/mL	5 mmol	10 mL	Large vein push over 5 mins DO NOT mix with other drugs
Resonium A	–	0.25 g/kg	Mix 1 scoop (15 g) with 60 mL water	0.25 g/mL	1.25 g	5 mL	Oral, nasogastric or rectal
Hypocalcaemia – Critical (↓ calcium) Calcium gluconate (2.2 mmol/10 mL)	0.22 mmol/mL	0.11 mmol/kg	Undiluted	0.22 mmol/mL	0.55 mmol	2.5 mL	Large vein push over 3 - 5 mins DO NOT give with sodium bicarbonate
Hypomagnesaemia or Arrhythmia Magnesium Sulfate (10 mmol/5 mL)	2 mmol/mL	0.2 mmol/kg	Dilute 5 mL (10 mmol) to 50 mL	0.2 mmol/mL	1 mmol	5 mL	Pulse absent – Push over 3 - 5 mins Pulse present – Infuse over 20 mins

Trauma	Vial concentration	Recommended dose/kg	Dilution – Sodium Chloride 0.9%	Final concentration	Dose	Final volume	Administration
Blood – Initial		10 mL/kg			50 mL	50 mL	As clinically indicated
Tranexamic Acid – 1000 mg/10 mL	100 mg/mL	15 mg/kg	Dilute 10 mL (1000 mg) to 100 mL	10 mg/mL	75 mg	7.5 mL	Infuse over 10 mins

For ongoing bleeding refer to local Massive Haemorrhage Protocol for blood and product replacement

Analgesia	Vial concentration	Recommended dose/kg	Dilution – Sodium Chloride 0.9%	Final concentration	Dose	Final volume	Administration
Fentanyl – Nasal (100 microg/2 mL) Use Mucosal Atomiser Device (MAD)	50 microg/mL	1.5 microg /kg	Undiluted	50 microg/mL	7.5 microg	0.15 mL	Add 0.1 mL to initial dose to accommodate (MAD) dead space May repeat after 5 - 10 mins
Fentanyl – IV (100 microg/2 mL)	50 microg/mL	0.5 - 1 microg/kg	Dilute 2 mL (100 microg) to 10 mL	10 microg/mL	2.5 microg	0.25 mL	Dose may be repeated after 5 mins if required
Morphine – IV (10 mg/mL)	10 mg/mL	0.05 - 0.1 mg/kg	Dilute 1 mL (10 mg) to 10 mL	1 mg/mL	0.25 mg	0.25 mL	Dose may be repeated after 5 mins if required

5kg

Analgesia if intraosseous IO drug or fluid administration causes pain	Vial concentration	Recommended dose/kg	Preparation	Final concentration	Dose	Final volume to administer	Administration
Lignocaine (Lidocaine) 1% IO	10 mg/mL (1%)	0.5 mg/kg	Undiluted	10 mg/mL	2.5 mg	0.25 mL	Instil dose - Follow with 1 mL slow push of sodium chloride 0.9% over 1-2 minutes. Allow to dwell for 1 minute. Rapid flush with 5 mL. Half original dose can be repeated as above

Antiarrhythmics - only in consultation with a Paediatric Cardiologist	Vial concentration	Recommended dose/kg	Preparation	Final concentration	Dose	Final volume	Administration
AmiODARone (Load) 150 mg/3 mL	See Infusion guide for doses and administration directions						
Esmolol 100 mg/10 mL	10 mg/mL	0.25 - 0.5 mg/kg	Undiluted	10 mg/mL	1.3 mg	0.13 mL	LOAD – Push over 1 - 2 mins. Continuous infusion may be considered after loading dose
Verapamil 5 mg/2 mL	2.5 mg/mL	0.1 mg/kg	Dilute 2 mL (5 mg) up to 10 mL	0.5 mg/mL	Consult	Consult	Infuse over 5 - 10 mins

Acute behavioural disturbance Oral	Medication preparation	Recommended dose	Preparation	Final concentration	Dose	Final volume	Administration
Diazepam	Liquid 1 mg/mL 5 mg tablet	0.2 mg/kg	If liquid unavailable dissolve 5 mg tablet in 5 mL of water	1 mg/mL	Consult	Consult	If tablet preferred round dose to nearest half or full tablet
Lorazepam	1 mg tablet	0.5 - 1 mg	Tablet may be dissolved in small volume of water		Consult	Consult	
Olanzapine	2.5 - 5 mg wafer	2.5 - 5 mg	Place wafer on top of tongue		Consult	Consult	Max daily dose not to exceed 20 mg
Risperidone	1 mg/mL liquid or tablets	0.02 - 0.04 mg/kg	If liquid unavailable tablets may be dissolved to make 1 mg/mL solution	1 mg/mL	Consult	Consult	Dose has been rounded

Acute behavioural disturbance IM	Vial concentration	Recommended dose	Preparation	Final concentration	Dose	Final volume	Administration
Droperidol	2.5 mg/mL	0.1 - 0.2 mg/kg	Undiluted	2.5 mg/mL	Consult	Consult	Max single dose not to exceed 10 mg. Total daily dose 0.4 mg/kg not to exceed 20 mg
Olanzapine	10 mg/mL	2.5 - 5 mg	Reconstitute vial with 2.1 mL WFI	5 mg/mL	Consult	Consult	Max single dose not to exceed 10 mg. Total daily dose 0.4 mg/kg not to exceed 20 mg

Reversal dystonia	Vial concentration	Recommended dose/kg	Preparation	Final concentration	Dose	Final volume	Administration
Benztropine (Benzatropine) IV or IM 2 mg/2 mL	1 mg/mL	0.02 mg/kg	Undiluted	1 mg/mL	Consult	Consult	IV or IM

5kg

Drug	Vial concentration	Recommended dose/kg range	Preparation		Final rate range	Administration/ route
			Glucose 5% or Sodium Chloride 0.9%	Final concentration		
Open Ductus Arteriosus						
Alprostadil (Prostaglandin/PGE1)	500 microg/mL	50 to 100 nanogram/kg/min	Dilute 0.2 mL (100 microg) to 50 mL	2 microg/mL (2000 nanogram/mL)	7.5 to 15 mL/hr	IV

Inotropes						
Adrenaline (Epinephrine)	1:1000; 1 mg/mL	0.05 to 1 microg/kg/min	Dilute 1 mL (1 mg) to 50 mL	20 microg/mL	0.8 to 15 mL/hr	IV
Dobutamine	250 mg/20 mL	2 to 20 microg/kg/min	Dilute 6 mL (75 mg) to 50 mL	1.5 mg/mL	0.4 to 4 mL/hr	IV
Dopamine	200 mg/5 mL	2 to 20 microg/kg/min	Dilute 1.5 mL (60 mg) to 50 mL	1.2 mg/mL	0.5 to 5 mL/hr	IV
Noradrenaline (Norepinephrine)	4 mg/4 mL	0.05 to 1 microg/kg/min	Dilute 1 mL (1 mg) to 50 mL	20 microg/mL	0.8 to 15 mL/hr	IV

Antiarrhythmics - only in consultation with a Paediatric Cardiologist						
AmiODAROne <u>LOAD</u>	50 mg/mL	25 microg/kg/min (for 4 hrs)	Dilute 2 mL (100 mg) to 50 mL in Glucose 5%	2 mg/mL	Dose 30 mg (15 mL) infuse at 3.8 mL/hr	IV
AmiODAROne [after loading dose]	50 mg/mL	5 to 15 microg/kg/min	Dilute 2 mL (100 mg) to 50 mL in Glucose 5%	2 mg/mL	0.8 to 2.3 mL/hr	IV
Esmolol	100 mg/10 mL	50 to 200 microg/kg/min	Undiluted – draw up 50 mL (500 mg)	10 mg/mL	1.5 to 6 mL/hr	IV

Sedation						
Fentanyl	100 microg/2 mL	1 to 10 microg/kg/hr	Dilute 10 mL (500 microg) to 50 mL	10 microg/mL	0.5 to 5 mL/hr	IV
Midazolam	Various strengths	30 to 120 microg/kg/hr	Dilute 10 mg to 50 mL	0.2 mg/mL	0.8 to 3 mL/hr	IV
Morphine	Various strengths	5 to 80 microg/kg/hr	Dilute 5 mg to 50 mL	0.1 mg/mL	0.3 to 4 mL/hr	IV

Paralytic Agents – only on discussion with Paediatric Intensivist						
Vecuronium	10 mg vial	1 to 3 microg/kg/min	Dilute 25 mL (50 mg) to 50 mL	1 mg/mL	0.3 to 0.9 mL/hr	IV

Electrolytes						
Hyperkalaemia Glucose 10%	–	5 mL/kg/hr	Use a glucose 10% bag - Undiluted <i>Administer with Actrapid infusion</i>	10%	25 mL/hr	IV. Run insulin and glucose infusions (concurrently) until K+ within range monitor BSLs
AND ACTRAPID (Insulin neutral)	300 units/3 mL	0.1 units/kg/hr	Dilute 0.5 mL (50 units) to 50 mL with Sodium Chloride 0.9% <i>Administer with Glucose infusion</i>	1 unit/mL	0.5 mL/hr	